



Soils Map



Soils data provided by USDA and NRCS.



State: **South Dakota**
 County: **Tripp**
 Location: **29-98N-77W**
 Township: **Weaver**
 Acres: **640**
 Date: **10/1/2018**



Maps Provided By:



Area Symbol: SD123, Soil Area Version: 18

| Code | Soil Description | Acres | Percent of field | PI Legend | Non-Irr Class *c | Irr Class *c | Productivity Index | State Productivity Index | NCCPI Soybeans |
|-------------------------|--|--------|------------------|-----------|------------------|--------------|--------------------|--------------------------|----------------|
| BOD | Boyd-Okaton association, 9 to 25 percent slopes | 285.13 | 44.6% | | Vle | | 16 | 11 | 15 |
| BnC | Boyd clay, 5 to 9 percent slopes | 90.86 | 14.2% | | IVe | | 49 | 33 | 20 |
| MfE | Manter-Anselmo fine sandy loams, 15 to 30 percent slopes | 43.68 | 6.8% | | Vle | Vle | 19 | 13 | 27 |
| PsA | Wendte clay, channeled, 0 to 2 percent slopes | 36.44 | 5.7% | | VIw | | 34 | 23 | 25 |
| MaB | Manter fine sandy loam, 3 to 9 percent slopes | 35.45 | 5.5% | | IIIe | IIIe | 52 | 35 | 41 |
| RaB | Ree loam, 3 to 6 percent slopes | 31.03 | 4.8% | | Ile | | 79 | 54 | 56 |
| AbB | Anselmo fine sandy loam, 3 to 6 percent slopes | 28.12 | 4.4% | | IIIe | IIIe | 54 | 37 | 34 |
| DgB | Doger loamy fine sand, 0 to 6 percent slopes | 26.56 | 4.1% | | IVe | | 32 | 22 | 25 |
| RaC | Ree loam, 6 to 9 percent slopes | 18.34 | 2.9% | | IIIe | Ile | 72 | 49 | 48 |
| WgB | Wewela fine sandy loam, 3 to 6 percent slopes | 15.27 | 2.4% | | IVe | IVe | 39 | 27 | 22 |
| MoB | Millboro silty clay, 3 to 6 percent slopes | 15.23 | 2.4% | | IIIe | IIIe | 79 | 54 | 28 |
| AaB2 | Anselmo loamy fine sand, 0 to 9 percent slopes, eroded | 12.29 | 1.9% | | Vle | | 47 | 32 | 35 |
| W | Water | 1.60 | 0.2% | | | | 0 | 0 | 0 |
| Weighted Average | | | | | | | 33.5 | 22.8 | 23.5 |

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.